arXiv:1210.3617v1 [gr-qc] 12 Oct 2012

Irreversible thermodynamic description of interacting dark energy - dark matter cosmological models

Tiberiu Harko[1]* and Francisco S.N. Lobo[2]†
[1]*Department of Physics and Center for Theoretical and Computational Physics,*
The University of Hong Kong, Pok Fu Lam Road, Hong Kong and
[2]*Centro de Astronomia e Astrofísica da Universidade de Lisboa,*
Campo Grande, Ed. C8 1749-016 Lisboa, Portugal

(Dated: October 16, 2012)

We investigate the interaction between dark energy and dark matter in the framework of irreversible thermodynamics of open systems with matter creation/annihilation. We consider dark energy and dark matter as an interacting two component (scalar field and "ordinary" dark matter) cosmological fluid in a homogeneous spatially flat and isotropic Friedmann-Robertson-Walker (FRW) Universe. The thermodynamics of open systems as applied together with the gravitational field equations to the two component cosmological fluid leads to a generalisation of the elementary dark energy-dark mater interaction theory, in which the decay (creation) pressures are explicitly considered as parts of the cosmological fluid stress-energy tensor. Specific models describing coherently oscillating scalar waves, leading to a high particle production at the beginning of the oscillatory period, and models with a constant potential energy scalar field are considered. Furthermore, exact and numerical solutions of the gravitational field equations with dark energy-dark matter interaction are also obtained.

PACS numbers: 04.50.Kd, 04.20.Cv, 04.20.Fy

I. INTRODUCTION

The observational data from type Ia supernovae, initially reported in [1], have generated a large theoretical and observational effort for the understanding of the observed present accelerated expansion of the Universe. Subsequent work on type Ia supernovae [2], the cosmic microwave background [3], and baryon acoustic oscillations [4] fully support the initial interpretation of the observational data that the expansion of the Universe is accelerating. The late-time cosmic acceleration is usually assumed to be driven by a fluid/field generically denoted dark energy [5]. Presently very little is known about dark energy, namely, its possible composition or its structure. Two main scenarios have been proposed to explain the nature of the dark energy: a cosmological constant Λ [5], or a scalar field, usually called quintessence [6]. The action for gravity and the scalar field is $S = \int \left[R/16\pi G - (1/2)\nabla^{\alpha}\phi\nabla_{\alpha}\phi - V(\phi)\right]\sqrt{-g}d^4x$, where $V(\phi)$ is the self-interaction potential [7].

Another of the central issues in modern astrophysics is the dark matter problem (see [8] for an extensive review of the recent results of the search for dark matter). The necessity of considering the existence of dark matter at a galactic and extragalactic scale is required by two fundamental observational evidences: the behavior of the galactic rotation curves, and the mass discrepancy in clusters of galaxies, respectively. On the galactic/intergalactic scale the rotation curves of spiral galaxies [9–11] provide compelling evidences pointing towards the problems Newtonian gravity and/or standard general relativity has to face at these scales. The behavior of the galactic rotation curves and of the virial mass of galaxy clusters is usually explained by postulating the existence of some dark (invisible) matter, distributed in a spherical halo around the galaxies. The dark matter is assumed to be a cold, pressure-less medium. Many possible candidates for dark matter have been proposed, the most popular ones being the WIMPs (Weakly Interacting Massive Particles) (for a review of the particle physics aspects of the dark matter see [12]). While extremely small, their interaction cross sections with normal baryonic matter, are expected to be non-zero, so that their direct experimental detection may be possible.

In this context, cosmological evolution and dynamics are largely dominated by dark energy and dark matter. Dark energy has a repulsive effect, driving the Universe to accelerate, while dark matter is gravitationally attractive. In the standard approach to cosmology there is no interaction between these two components. Since the gravitational effects of the dark energy and of the dark matter are opposite (i.e., gravitational repulsion versus gravitational attraction) and since dark energy is very homogeneously distributed, while dark matter clumps around ordinary matter, one expects that any dynamic interaction between these two dark components of the Universe would be extremely weak, or even negligible. However, the possibility of such an interaction cannot be excluded *a priori* and, following some early proposals [13], presently interacting dark matter-dark energy models were extensively investigated in the literature [14]. In the standard approach, one may model dark energy as a scalar field with energy density ρ_ϕ and pressure p_ϕ, while dark matter is described as a matter

---

*Electronic address: harko@hkucc.hku.hk
†Electronic address: flobo@cii.fc.ul.pt

fluid with density ρ_{DM} and pressure p_{DM}, satisfying an equation of state $w_{DM} = p_{DM}/\rho_{DM} \equiv 0$. By assuming a spatially flat Friedman-Robertson-Walker (FRW) background with scale factor, $a(t)$, and allowing for creation/annihilation between the dark energy (scalar field) and the dark matter fluid at a rate Q, the equations describing the variations of the dark energy and dark matter densities ρ_ϕ and ρ_{DM} can be written as [13, 14]

$$\dot{\rho_\phi} + 3H\left(1 + w_\phi\right)\rho_\phi = -Q, \tag{1}$$

$$\dot{\rho_{DM}} + 3H\left(1 + w_{DM}\right)\rho_{DM} = +Q, \tag{2}$$

respectively, where $H \equiv \dot{a}/a$ is the Hubble function. The derivatives with respect to the cosmological time, t, will be indicated in the following by an overdot. The dark energy and the dark matter create/decay into one another via the common creation/annihilation rate $\pm Q$. Hence Q describes the interaction between the two dark components of the Universe. When $Q > 0$, dark energy is converted to dark matter, while if $Q < 0$, dark matter is converted to dark energy. Since there is no fundamental theoretical approach that may specify the functional form of the coupling between dark energy and dark matter, presently coupling models are necessarily phenomenological, although one might view some couplings more physical or more natural than others. Hence a large number of functional forms for Q have been proposed, and investigated in the literature, such as $Q = \rho^0_{crit}(1+z)^3 H(z) I_Q(z)$, where z is the redshift, and $I_Q(z)$ is an interaction function that depends on the redshift [15], or $Q \propto \rho_{DM}\dot{\phi}$, and $Q \propto H\rho_{DM}$, respectively [16].

In writing down Eqs. (1) and (2), one assumes the idealised picture of a very quick decay in which the final products reach equilibrium states immediately. However, the dark energy/dark matter creation/annihilation period may be characterised by complicated nonequilibrium processes, with a highly nonequilibrium distribution of the produced particles, subsequently relaxing to an equilibrium state. Thermodynamical systems in which matter creation occurs fits in the class of open thermodynamical systems in which the usual adiabatic conservation laws are modified thereby including irreversible matter creation [17]. The thermodynamics of open systems were first applied to cosmology in [17]. Explicit inclusion of the matter creation in the matter stress-energy tensor in the Einstein field equations leads to a three stage cosmological history, starting from an instability of the vacuum. During the first stage the Universe is driven from an initial fluctuation of the vacuum to a de Sitter phase, and particle creation occur. The de Sitter phase does exist during the decay time of its constituents (second stage) and ends, after a phase transition into the usual FRW Universe. The phenomenological approach of [17] was further discussed and generalised in [18] through a covariant formulation allowing specific entropy variation as usually expected for non-equilibrium processes. Cosmological models involving irreversible matter creation have been considered in [19].

It is the purpose of the present paper to apply the thermodynamics of open systems as developed in [17] and [18] to a cosmological fluid mixture consisting of two components: dark energy, described by a scalar field, and dark matter, modeled as an ordinary matter fluid, in which particle decay and production occur. This situation may be specific to both early and late stages of cosmological evolution. The thermodynamics of irreversible processes as applied to cosmological models with interacting dark energy and dark matter leads to a self-consistent description of the dark energy and dark matter particle creation/annihilation processes, which in turn determine the whole dynamics and future evolution of the Universe.

The present paper is organised as follows. In Section II we present, in some detail, due to its important role, the thermodynamical theory of irreversible matter creation processes. The theory is applied to a two-component cosmological fluid with interacting dark energy and dark matter, and the resulting gravitational field equations are written down in Section III. Particular models, and exact and numerical solutions to the field equations are considered in Section IV. In Section V we discuss and conclude our results. Throughout the paper we use a system of units so that $8\pi G = c = 1$.

II. THERMODYNAMICS OF IRREVERSIBLE COSMOLOGICAL MATTER CREATION

We consider a cosmological volume element V containing N particles. For a closed system, N is constant, and the corresponding thermodynamic conservation of the internal energy E is expressed by the first law of thermodynamics as [17]

$$dE = dQ - pdV, \tag{3}$$

where dQ is the heat received by the system during time dt, p is the thermodynamic pressure, and V is any comoving volume. By introducing the energy density ρ defined as $\rho = E/V$, the particle number density n given by $n = N/V$, and the heat per unit particle dq, with $dq = dQ/N$, Eq. (3) becomes

$$d\left(\frac{\rho}{n}\right) = dq - pd\left(\frac{1}{n}\right). \tag{4}$$

Equation (4) is also valid for open systems in which N is time dependent, $N = N(t)$. Equation (3) must be modified to explicitly take into account the variation of the particle number,

$$d\left(\rho V\right) = dQ - pdV + \left(\frac{h}{n}\right) d\left(nV\right), \tag{5}$$

where $h = \rho + p$ is the enthalpy (per unit volume) of the fluid. For open systems, adiabatic transformations ($dQ = 0$) are defined by the equation

$$d(\rho V) + pdV - \left(\frac{h}{n}\right) d(nV) = 0. \tag{6}$$

Equation (6) can be written in a number of equivalent forms as

$$\dot{\rho} = \left(\frac{h}{n}\right)\dot{n}, \tag{7}$$

or, equivalently,

$$p = \dot{\rho} - \rho\frac{\dot{n}}{n}. \tag{8}$$

The Einstein field equations

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = T_{\mu\nu}, \tag{9}$$

involve the macroscopic stress-energy tensor $T_{\mu\nu}$, which, in the cosmological case, corresponds to a perfect fluid. It is characterised by a phenomenological energy density ρ and pressure $\bar{p}$, and its components are given by

$$T_0^0 = \rho, T_1^1 = T_2^2 = T_3^3 = -\bar{p}. \tag{10}$$

In addition to the Einstein field equations we have the Bianchi identities, which lead to $\nabla_\nu T_\mu^\nu = 0$, and to the relation

$$d(\rho V) = -\bar{p}dV. \tag{11}$$

In the presence of adiabatic irreversible matter creation the appropriate analysis must be performed in the context of open systems. This involves the inclusion of a supplementary creation/annihilation pressure p_c, as we may write Eq. (6) in a form similar to Eq. (11), namely [17]

$$d(\rho V) = -\left(p + p_c\right)dV, \tag{12}$$

where the creation pressure is given by

$$p_c = -\left(\frac{h}{n}\right)\frac{d(nV)}{dV} = -\left(\frac{h}{n}\right)\frac{V}{\dot{V}}\left(\dot{n} + \frac{\dot{V}}{V}n\right). \tag{13}$$

Creation of matter corresponds to a (negative) supplementary pressure p_c, which must be considered as part of the cosmological pressure $\bar{p}$ entering into the Einstein field equations (decaying of matter leads to a positive decay pressure),

$$\bar{p} = p + p_c. \tag{14}$$

The entropy change dS in an open thermodynamic system can be decomposed into an entropy flow d_0S, and the entropy creation d_iS,

$$dS = d_0S + d_iS, \tag{15}$$

with $d_iS \geq 0$. To evaluate dS we start from the total differential of the entropy,

$$Td(sV) = d(\rho V) + pdV - \mu d(nV), \tag{16}$$

where $s = S/V \geq 0$ and $\mu n = h - Ts$, $\mu \geq 0$ being the chemical potential. In a homogeneous system $d_0S = 0$, but matter creation contributes to the entropy production. From Eqs. (15) and (16) we obtain [17]

$$T\frac{dS}{dt} = T\frac{d_iS}{dt} = T\frac{s}{n}\frac{d(nV)}{dt}. \tag{17}$$

To complete the problem we need one more relation between the particle number n and V, describing the time dynamics of n as a result of matter creation (decay) processes. This relation is usually taken in the form [17, 18]

$$\frac{1}{V}\frac{d(nV)}{dt} = \Psi(t), \tag{18}$$

where $\Psi(t)$ is the matter creation (or decay) rate ($\Psi(t) > 0$ corresponds to particle creation, while $\Psi(t) < 0$ corresponds to particle decay). The creation pressure (13) depends on the matter creation (decay) rate, thereby coupling Eqs. (13) and (18) to each other and, although indirectly, both of them with the energy conservation law (12), which is contained in the Einstein field equations themselves. The entropy production can also be expressed as a function of the matter creation rate as

$$S(t) = S\left(t_0\right)e^{\int_{t_0}^t \frac{\Psi(t)}{n}dt}. \tag{19}$$

III. COSMOLOGICAL DYNAMICS IN A UNIVERSE WITH IRREVERSIBLE DARK ENERGY-DARK MATTER INTERACTION

We shall model the Universe as an open thermodynamical system, consisting of a two-component (dark energy and dark matter) perfect fluid, with the particle number densities denoted by n_ϕ, and n_{DM}, respectively. n_ϕ corresponds to the "particles" of the scalar field, while n_{DM} is the particle number of the dark matter. We denote the corresponding energy densities by ρ_ϕ and ρ_{DM}, respectively. The stress-energy tensor of the two-component cosmological fluid is given by

$$T_\mu^\nu = T_\mu^{(\phi)\nu} + T_\mu^{(DM)\nu} = \rho u_\mu u^\nu - \bar{p}\,\delta_\mu^\nu, \tag{20}$$

where $u^\mu = dx^\mu/ds$ is the four-velocity, and

$$\rho = \rho_\phi + \rho_{DM}, \qquad \bar{p} = \bar{p}_\phi + \bar{p}_{DM}. \tag{21}$$

The energy density and pressure of the dark energy are given by $\rho_\phi = \dot{\phi}^2/2 + U(\phi)$ and $p_\phi = \dot{\phi}^2/2 - U(\phi)$, respectively, where $U(\phi)$ is the self-interaction potential. We suppose that neither the particle numbers nor the stress-energy of the components are separately conserved, that is, particle inter-conversion and exchange of energy and momentum between the two components are admitted. The cosmological fluid mixture is characterised by a total energy density $\rho = \rho_\phi + \rho_{DM}$, total thermodynamic pressure $\bar{p} = \bar{p}_\phi + \bar{p}_{DM}$ and a total particle number $n = n_\phi + n_{DM}$. We consider that the geometry of the spacetime is described by the flat FRW line element, given by

$$ds^2 = dt^2 - a^2(t)\left(dx^2 + dy^2 + dz^2\right). \tag{22}$$

We shall assume that the particle number densities n_ϕ and n_{DM} of each component of the fluid obey the following balance laws,

$$\dot{n}_\phi + 3Hn_\phi = -\Gamma_1\rho_\phi, \tag{23}$$
$$\dot{n}_{DM} + 3Hn_{DM} = \Gamma_2\rho_\phi, \tag{24}$$

respectively, where $\Gamma_1 \neq 0$ and $\Gamma_2 \neq 0$ are arbitrary functions. Equations (23) and (24) describe the decay of the dark energy ϕ-particles, and the creation of the dark matter particles, with a scalar field decay rate and a dark matter creation rate $\Psi(t) \propto \rho_\phi$. Thus, the dynamics of ϕ-particle decay and the creation of dark matter is governed in the present model by the scalar field via its energy density. From Eqs. (23) and (24) it follows that the total particle number n obeys the balance equation

$$\dot{n} + 3Hn = \left(\Gamma_2 - \Gamma_1\right)\rho_\phi. \tag{25}$$

Hence in the case of an interacting dark matter and dark energy the total particle number conservation occurs only in very special cases, and therefore we shall suppose that generally $\Gamma_1 \neq \Gamma_2$.

In the framework of the thermodynamics of irreversible processes particle creation and decay gives rise to a decay and a creation thermodynamic pressure, given by

$$p_c^{(\phi)} = \frac{\Gamma_1\left(\rho_\phi + p_\phi\right)\rho_\phi}{3Hn_\phi}, \tag{26}$$

and

$$p_c^{(DM)} = -\frac{\Gamma_2\left(\rho_{DM} + p_{DM}\right)\rho_\phi}{3Hn_{DM}}, \tag{27}$$

respectively, while the total creation pressure becomes

$$\begin{aligned} p_c^{(total)} &= p_c^{(\phi)} + p_c^{(DM)} \\ &= \frac{\rho_\phi}{3H}\left[\frac{\Gamma_1\left(\rho_\phi + p_\phi\right)}{n_\phi} - \frac{\Gamma_2\left(\rho_{DM} + p_{DM}\right)}{n_{DM}}\right] \end{aligned} \tag{28}$$

Using the results obtained above the complete Einstein gravitational field equations describing the dynamics of a flat FRW spacetime filled with a mixture of interacting dark matter (scalar field) and dark matter can be expressed in the form

$$3H^2 = \rho_\phi + \rho_{DM}, \tag{29}$$

$$\begin{aligned} 2\dot{H} + 3H^2 &= -p_\phi - p_{DM} \\ &- \frac{\rho_\phi}{3H}\left[\frac{\Gamma_1\left(\rho_\phi + p_\phi\right)}{n_\phi} - \frac{\Gamma_2\left(\rho_{DM} + p_{DM}\right)}{n_{DM}}\right], \end{aligned} \tag{30}$$

and

$$\dot{n}_\phi + 3Hn_\phi = -\Gamma_1\rho_\phi, \tag{31}$$
$$\dot{n}_{DM} + 3Hn_{DM} = \Gamma_2\rho_\phi, \tag{32}$$

where $\rho_{DM} = \rho_{DM}\left(n_{DM}\right)$ and $p_{DM} = p_{DM}\left(n_{DM}\right)$, and recalling that the energy density and pressure of dark energy is given by

$$\rho_\phi = \frac{\dot{\phi}^2}{2} + U(\phi), \qquad p_\phi = \frac{\dot{\phi}^2}{2} - U(\phi). \tag{33}$$

As applied to each component of the cosmological fluid, Eq. (7), the second law of thermodynamics for open systems provides the following relationships

$$\dot{\rho}_\phi + 3H\left(\rho_\phi + p_\phi\right) + \frac{\Gamma_1\left(\rho_\phi + p_\phi\right)\rho_\phi}{n_\phi} = 0, \tag{34}$$

$$\dot{\rho}_{DM} + 3H\left(\rho_{DM} + p_{DM}\right) = \frac{\Gamma_2\left(\rho_{DM} + p_{DM}\right)\rho_\phi}{n_{DM}}, \tag{35}$$

respectively.

Equation (34), which describes the dynamics of the dark energy during its interaction with dark matter, can be written in an equivalent form as

$$\ddot{\phi} + 3H\dot{\phi} + \Gamma\left(\phi, \dot{\phi}, U\right)\dot{\phi} + U'(\phi) = 0, \tag{36}$$

where we have denoted $\Gamma\left(\phi, \dot{\phi}, U\right) = \Gamma_1\rho_\phi/n_\phi$. Therefore in the framework of the thermodynamics of irreversible processes a friction term in the scalar field Eq. (36) arises naturally, and in a general form, as a direct consequence of the second law of thermodynamics as applied to an open system.

Adding Eqs. (34) and (35), the evolution of the total energy density $\rho = \rho_\phi + \rho_{DM}$ of the cosmological fluid is governed by the equation

$$\begin{aligned} &\dot{\rho} + 3H\left(\rho + p_\phi + p_{DM}\right) = \\ &\rho_\phi\left[\frac{\Gamma_2\left(\rho_{DM} + p_{DM}\right)}{n_{DM}} - \frac{\Gamma_1\left(\rho_\phi + p_\phi\right)}{n_\phi}\right]. \end{aligned} \tag{37}$$

For the entropy of the newly created matter we obtain

$$S_{DM}(t) = S_{DM}\left(t_0\right)\exp\left(\int_{t_0}^{t}\Gamma_2\frac{\rho_\phi}{n_{DM}}dt\right). \tag{38}$$

Irreversible matter particle creation is an adiabatic process, the produced entropy being entirely due to the increase in the number of fluid particles, there being no increase in the entropy per particle due to dissipative processes.

IV. IRREVERSIBLE DARK ENERGY-DARK MATTER INTERACTION MODELS

In the present Section we consider, within the framework of irreversible thermodynamics with matter creation/annihilation, a number of specific cosmological models with dark energy-dark matter interaction. As a first case we consider the situation in which the density of the dark matter is much smaller than the energy density of the scalar field. This case corresponds to an Universe dominated by the dark energy component, assumed to be represented by a coherent wave of ϕ-particles. In this case the kinetic term dominates in the total energy of the scalar field. As a second case we consider a potential energy dominated scalar field. The general dynamics of the cosmological model with interacting dark energy and dark matter is also considered, and the cosmological evolution equations are studied numerically.

A. Coherent Scalar Waves-Dark matter interaction

We shall consider in the following that the energy density and particle number of the newly created dark matter is much smaller than the energy density and particle number of the corresponding scalar field fluid component, that is the relations $\rho_{DM} \ll \rho_\phi$, $n_{DM} \ll n_\phi$, and $p_{DM} \ll p_\phi$ hold. In this case the Universe is dominated by the scalar field energy density, and its evolution is not influenced by the matter content. We shall work throughout with finite values of the fluid quantities at $t = t_0$. The coupling between scalar field and dark matter is realised only by means of the balance equation of ordinary matter via the scalar field energy density, and the basic equations describing the dynamics of a flat FRW scalar field filled space-time interacting with a dark matter component are given by

$$
\begin{aligned}
3H^2 &= \rho_\phi \qquad (39)\\
2\dot{H} + 3H^2 &= -p_\phi - \frac{\Gamma_1\left(\rho_\phi + p_\phi\right)\rho_\phi}{3Hn_\phi}, \qquad (40)
\end{aligned}
$$

and

$$
\dot{\rho}_\phi = \left(\rho_\phi + p_\phi\right)\frac{\dot{n}_\phi}{n_\phi}, \qquad (41)
$$

$$
\begin{aligned}
\dot{n}_\phi + 3Hn_\phi &= -\Gamma_1\rho_\phi, \qquad (42)\\
\dot{n}_{DM} + 3Hn_{DM} &= \Gamma_2\rho_\phi, \qquad (43)
\end{aligned}
$$

respectively.

A homogeneous scalar field oscillating with frequency m_ϕ can be considered as a coherent wave of "particles" with zero momenta, and with a particle number density given by [20]

$$
n_\phi = \frac{\rho_\phi}{m_\phi}, \qquad m_\phi = \text{constant}. \qquad (44)
$$

In other words, n_ϕ oscillators of the same frequency m_ϕ oscillating coherently with the same phase can be described as a single homogeneous wave $\phi(t)$. Insertion of the energy density of the scalar field given by Eq. (44) in Eq. (41) leads to the condition

$$
p_\phi = 0, \qquad (45)
$$

or, equivalently,

$$
U\left(\dot{\phi}\right) = \frac{\dot{\phi}^2}{2}. \qquad (46)
$$

Therefore, a homogeneous oscillating scalar field is described in the present model by a Barrow-Saich type potential, with potential energy of the scalar field proportional to the kinetic one [21]. The energy density of the scalar field becomes $\rho_\phi = \dot{\phi}^2$, and this relation, obtained naturally in the framework of the present formalism is very similar to the equation $\rho_\phi = \left\langle\dot{\phi}^2\right\rangle$ obtained by replacing $\dot{\phi}^2$ by its average value per cycle [22].

In this case the equations describing the dynamics of the FRW type spacetime filled by the decaying oscillating homogeneous scalar field in the presence of dark matter creation become

$$
3H^2 = \dot{\phi}^2, \qquad (47)
$$

$$
2\dot{H} + 3H^2 = -\Gamma_1 m_\phi H, \qquad (48)
$$

$$
\dot{n}_{DM} + 3Hn_{DM} = 3\Gamma_2 H^2, \qquad (49)
$$

respectively. In the following we will assume, for simplicity, that Γ_1 and Γ_2 are constants. By introducing a set of dimensionless variables τ, h and θ_{DM} by means of the transformations

$$
t = \frac{2}{\Gamma_1 m_\phi}\tau, \quad H = \frac{\Gamma_1 m_\phi}{3}h, \quad n_{DM} = \frac{2\Gamma_1\Gamma_2 m_\phi}{3}\theta_{DM}, \qquad (50)
$$

Equations (48) and (49) take the form

$$
\frac{dh}{d\tau} = -h(h+1), \qquad (51)
$$

$$
\frac{d\theta_{DM}}{d\tau} + 2h\theta_{DM} = h^2, \qquad (52)
$$

and yield the following general solutions

$$
h(\tau) = \frac{1}{e^\tau - 1}, \qquad (53)
$$

and

$$
\theta_{DM} = \frac{\left[\left(e^{\tau_0} - 1\right)^2\theta_{DM0} + 1/2\right]e^{2(\tau-\tau_0)} - 1/2}{\left(e^\tau - 1\right)^2}, \qquad (54)
$$

respectively, where we have denoted $\theta_{DM0} = \theta_{DM}\left(\tau_0\right)$.

The evolution of the scale factor is given by

$$
a(\tau) = a_0\left(\frac{e^\tau - 1}{e^\tau}\right)^{2/3}. \qquad (55)
$$

The deceleration parameter $q = d(1/H)/dt - 1$ is given by

$$
q = \frac{3}{2}e^\tau - 1. \qquad (56)
$$

For $(t-t_0) \ll \Gamma_1^{-1}$, i.e., at the start of the oscillatory period corresponding to the dark matter production, the approximate solution of the field equations is given by

$$h \approx \frac{1}{\tau}, \qquad a \approx a_0 \tau^{2/3}, \tag{57}$$

$$\rho_\phi \approx \frac{1}{\tau^2}, \qquad \theta_{DM} \approx \theta_{DM0}\frac{\tau_0^2}{\tau^2}\left(1+\frac{\tau-\tau_0}{\theta_{DM0}\tau_0^2}\right). \tag{58}$$

This phase corresponds to an Einstein-de Sitter expansion, with decaying dark energy, and dark matter creation.

During the initial oscillating period of the scalar field dominated FRW Universe there is a rapid increase of its dark matter content. The particle number density increases during a time interval

$$\Delta t = t_{max} - t_0 = \left(1-\frac{3\theta_{DM0}t_0}{2\Gamma_2}\right)t_0, \tag{59}$$

and reaches a maximum value given by

$$n_{DM}^{(max)} = \frac{\Gamma_2}{3\Delta t}\frac{1-3\theta_{DM0}t_0/2\Gamma_2}{t_0\left(1-3\theta_{DM0}t_0/4\Gamma_2\right)}. \tag{60}$$

Assuming that the scalar field oscillations decay into relativistic dark matter, the energy density and temperature of the dark matter component of the cosmological fluid is given by $\rho_{DM} \sim n^\gamma$, $T_{DM} \sim n^{\gamma-1}$, where $\gamma = 2$ corresponds to a stiff (Zeldovich) fluid obeying an equation of state of the form $\rho_{DM} = p_{DM}$, and $\gamma = 4/3$ corresponds to a relativistic, radiation like fluid.

The entropy produced during dark matter particle creation can be easily obtained from Eq. (38) and is given, in first approximation, by

$$\frac{S_{DM}(t)}{S_{DM0}} = \frac{\tau-\tau_0}{\theta_{DM0}\tau_0^2} + 1. \tag{61}$$

Therefore, for small times, there is a linear increase of the entropy of the dark energy dominated flat spacetime.

B. Constant scalar field potential dark energy and dark matter interaction

As a second example in the study of the irreversible interaction between dark energy (a scalar field) and dark matter we consider the case in which the scalar field potential may be approximated, at least for a certain time interval, as a positive constant, $U(\phi) = \Lambda = \text{constant} > 0$. Therefore, the energy density and the pressure of the scalar field can be written as

$$\rho_\phi = \frac{1}{2}\dot{\phi}^2 + \Lambda, \qquad \rho_\phi = \frac{1}{2}\dot{\phi}^2 - \Lambda. \tag{62}$$

Moreover, we assume that the change in the scalar field energy density due to the cosmological expansion can be neglected in the scalar field evolution equations, that is, in the energy density and particle number equations of the scalar field, given by Eqs. (31) and (34), the term containing $3H$ can be neglected. In this approximation the main contribution to the temporal dynamics of the scalar field is due to its decay into dark matter particles, and not to the cosmological expansion. On the other hand, the feedback of the newly created dark matter particles on the cosmological dynamics cannot be neglected. Within this approximation from Eq. (31), describing the scalar field particles decay, we obtain first

$$\rho_\phi = -\frac{1}{\Gamma_1}\dot{n}_\phi, \tag{63}$$

which gives the scalar field particle number density as a function of the energy of the scalar field. By substituting this expression of ρ_ϕ into Eq. (34) gives

$$\ddot{n}_\phi + \Gamma_1\frac{\dot{\phi}^2}{n_\phi}\dot{n}_\phi = 0. \tag{64}$$

Taking into account that

$$\dot{\phi}^2 = 2\left(\rho_\phi - \Lambda\right) = 2\left(-\frac{\dot{n}_\phi}{\Gamma_1} - \Lambda\right), \tag{65}$$

we obtain the equation describing the dynamics of the scalar field particles as

$$n_\phi \ddot{n}_\phi - 2n_\phi^2 - 2\Gamma_1\Lambda\dot{n}_\phi = 0. \tag{66}$$

By denoting $\dot{n}_\phi = u$, $\ddot{n}_\phi = u\,du/dn_\phi$, Eq. (66) becomes

$$n_\phi\frac{du}{dn_\phi} = 2\left(u + \Gamma_1\Lambda\right), \tag{67}$$

providing

$$\dot{n}_\phi = N_1 n_\phi^2 - \Gamma_1\Lambda, \tag{68}$$

where N_1 is an arbitrary integration constant, which can be determined from the initial condition $\dot{n}_\phi\left(t_0\right) = -\Gamma_1\rho_\phi\left(t_0\right) = -\Gamma_1\rho_{\phi 0}$ and $n_\phi\left(t_0\right) = n_{\phi 0}$ as

$$N_1 = \frac{\Gamma_1\left(\Lambda - \rho_{\phi 0}\right)}{n_{\phi 0}^2}. \tag{69}$$

Therefore the general solution of Eq. (66) is given by

$$n_\phi(t) = \sqrt{\frac{\Gamma_1\Lambda}{N_1}}\frac{1+N_2e^{2\alpha t}}{1-N_2e^{2\alpha t}}, \tag{70}$$

where $\alpha = \sqrt{\Gamma_1\Lambda N_1}$, and

$$N_2 = \frac{n_{\phi 0} - \sqrt{\Gamma_1\Lambda/N_1}}{n_{\phi 0} + \sqrt{\Gamma_1\Lambda/N_1}}e^{-2\alpha t_0}. \tag{71}$$

By assuming that the newly created dark matter particles are pressureless, $p_{DM} = 0$, and the energy density of the dark matter component is $\rho_{DM} = m_{DM}n_{DM}$, where

m_{DM} is the mass of the dark matter particle, then the evolution equation for the dark matter density can be written as

$$\begin{aligned}\dot{\rho}_{DM}+3H\rho_{DM} &= -\frac{\Gamma_2}{\Gamma_1}\dot{n}_{\phi} \\ &= \Gamma_2\Lambda-\frac{\Gamma_2\left(\Lambda-\rho_{\phi 0}\right)}{n_{\phi 0}^2}n_{\phi}^2, \quad (72)\end{aligned}$$

By neglecting the effects of the expansion of the Universe the dark matter energy increases as

$$\begin{aligned}\rho_{DM}(t) &\approx \Gamma_2\left(\Lambda-\sqrt{\frac{\Gamma_1\lambda}{N_1}}\frac{\Lambda-\rho_{\phi 0}}{n_{\phi 0}^2}\right)t \\ &-\frac{\Gamma_2\left(\Lambda-\rho_{\phi 0}\right)}{\alpha n_{\phi 0}^2}\sqrt{\frac{\Gamma_1\lambda}{N_1}}\frac{1}{N_2e^{2\alpha t}-1}. \quad (73)\end{aligned}$$

In the general case, in which the expansion of the Universe is also taken into account, the system of equations describing the irreversible dark energy-dark matter interaction are given by

$$3H^2=\rho_{\phi}+\rho_{DM}, \quad (74)$$

$$\dot{n}_{\phi}+3Hn_{\phi}=-\Gamma_1\rho_{\phi}, \quad (75)$$

$$\dot{\rho}_{\phi}+6H\left(\rho_{\phi}-\Lambda\right)+\frac{2\Gamma_1\left(\rho_{\phi}-\Lambda\right)\rho_{\phi}}{n_{\phi}}=0, \quad (76)$$

$$\dot{\rho}_{DM}+3H\rho_{DM}=\Gamma_1 m_{DM}\rho_{\phi}, \quad (77)$$

where, for simplicity, we have assumed $\Gamma_1=\Gamma_2=$ constant. By introducing a set of dimensionless variables $(r_{\phi},N_{\phi},r_{DM},\tau)$, defined as

$$\rho_{\phi}=\Lambda r_{\phi}, \quad n_{\phi}=\Lambda N_{\phi}, \quad (78)$$

$$\rho_{DM}=\Lambda m_{DM}r_{DM}, \quad t=\tau/\Gamma_1, \quad (79)$$

the field equations Eqs. (74)-(77) can be written in a dimensionless form as

$$\frac{1}{a}\frac{da}{d\tau}=\lambda\sqrt{r_{\phi}+m_{DM}r_{DM}}, \quad (80)$$

$$\frac{dN_{\phi}}{d\tau}+3\lambda\sqrt{r_{\phi}+m_{DM}r_{DM}}N_{\phi}=-r_{\phi}, \quad (81)$$

$$\frac{dr_{\phi}}{d\tau}+6\lambda\sqrt{r_{\phi}+m_{DM}r_{DM}}\left(r_{\phi}-1\right)+\frac{2r_{\phi}\left(r_{\phi}-1\right)}{N_{\phi}}=0, \quad (82)$$

$$\frac{dr_{DM}}{d\tau}+3\lambda\sqrt{r_{\phi}+m_{DM}r_{DM}}r_{DM}=r_{\phi}, \quad (83)$$

respectively, where

$$\lambda=\frac{\sqrt{\Lambda/3}}{\Gamma_1}. \quad (84)$$



FIG. 1: Time variation of the scale factor of the interacting dark energy-dark matter filled Universe, for different values of the parameter λ: $\lambda=0.001$ (solid curve), $\lambda 0.005$ (dotted curve), $\lambda=0.009$ (dashed curve), and $\lambda=0.012$ (long dashed curve). The initial conditions used for the numerical integration of the cosmological evolution equations are $a(0)=0.05$, $N_{\phi}(0)=10$, $r_{\phi}(0)=1.5$, and $r_{DM}(0)=0.01$, respectively. For the mass m_{DM} of the dark matter particle we have assumed the value $m=1$ MeV.



FIG. 2: Time variation of the scalar field particle number N_{ϕ} in the interacting dark energy-dark matter filled Universe, for different values of the parameter λ: $\lambda=0.001$ (solid curve), $\lambda 0.005$ (dotted curve), $\lambda=0.009$ (dashed curve), and $\lambda=0.012$ (long dashed curve). The initial conditions used for the numerical integration of the cosmological evolution equations are $a(0)=0.05$, $N_{\phi}(0)=10$, $r_{\phi}(0)=1.5$, and $r_{DM}(0)=0.01$, respectively. For the mass m_{DM} of the dark matter particle we have assumed the value $m=1$ MeV.

The dynamics of the interacting dark energy-dark matter system is determined by two control parameters, λ, and the mass m_{DM} of the dark matter particle. The time variations of the scale factor a, of the scalar field particle number N_{ϕ}, of the scalar field energy r_{ϕ}, as well as the dark matter energy density r_{DM} are represented, for different values of the parameter λ, and for $m_{DM}=1$, in Figs. 1-4.



FIG. 3: Time variation of the dimensionless scalar field energy r_ϕ in the interacting dark energy-dark matter filled Universe, for different values of the parameter λ: $\lambda = 0.001$ (solid curve), $\lambda 0.005$ (dotted curve), $\lambda = 0.009$ (dashed curve), and $\lambda = 0.012$ (long dashed curve). The initial conditions used for the numerical integration of the cosmological evolution equations are $a(0) = 0.05$, $N_\phi(0) = 10$, $r_\phi(0) = 1.5$, and $r_{DM}(0) = 0.01$, respectively. For the mass m_{DM} of the dark matter particle we have assumed the value $m = 1$ MeV.



FIG. 4: Time variation of the dimensionless dark matter energy r_{DM} in the interacting dark energy-dark matter filled Universe, for different values of the parameter λ: $\lambda = 0.001$ (solid curve), $\lambda 0.005$ (dotted curve), $\lambda = 0.009$ (dashed curve), and $\lambda = 0.012$ (long dashed curve). The initial conditions used for the numerical integration of the cosmological evolution equations are $a(0) = 0.05$, $N_\phi(0) = 10$, $r_\phi(0) = 1.5$, and $r_{DM}(0) = 0.01$, respectively. For the mass m_{DM} of the dark matter particle we have assumed the value $m = 1$ MeV.

As one can see from Fig. 1, the interacting dark energy-dark matter filled Universe is an expansionary state, with the rate of the expansion, and the scale factor evolution, strongly dependent of the dimensionless parameter λ, which depends on the ratio of the (constant) scalar field potential Λ, and of the scalar field decay rate Γ_1. Accelerated expansion can also be obtained in the framework of the present model, the creation pressure corresponding to the irreversible decay of the scalar field, and the matter creation, can drive the Universe into a de Sitter type phase. The scalar field particle number decays during the cosmological evolution, as shown in Fig. 2. The decay rate strongly depends on the numerical value of λ. The dimensionless energy of the scalar field r_ϕ, shown in Fig. 3, tends in the large time limit to the value 1, corresponding to $\rho_\phi = \Lambda$, and to a de Sitter expansion. This shows that the decay of the scalar field is determined and controlled by the kinetic energy term of the field $\dot{\phi}^2/2$, which is the source of the dark matter particles creation. When the energy and the pressure of the scalar field are dominated by the scalar field potential Λ, $\rho_\phi = -p_\phi = \Lambda$, then $\rho_\phi + p_\phi = 0$, and from Eq. (34) it follows that $\rho_\phi =$ constant, and the scalar field energy cannot be converted any more into other type of particles. The energy density of the dark matter particles, presented in Fig. 4, increases in time due to the decay of the ϕ-particles.

V. DISCUSSIONS AND FINAL REMARKS

In the present paper we have shown that the thermodynamics of the open systems is a valuable tool for describing the interacting dark energy - dark matter phases of the general relativistic cosmological models. As applied to a two component (scalar field and dark matter) cosmological model, the thermodynamics of irreversible processes provides a generalization of the elementary theory of dark energy-dark matter interaction, which envisages that during the dark energy dominated phase of the expansion of the Universe, when the expansion slows down, the energy stored in the zero mode oscillations of the scalar field transforms into particles via single particle decay. Thus the model presented in this paper gives a rigorous thermodynamical foundation, and a natural generalisation, to the theory of dark energy-dark matter interaction. Particle decay (creation) gives rise to a supplementary decay (creation) pressure which has to be included as a distinct part in the stress-energy tensor of the cosmological mixture. We have considered only particle creation in a scalar field (dark energy) dominated Universes, leading to a model in which particle production rate is extremely high at the beginning of the oscillatory period, and afterwards it tends to zero, when the scalar field energy and pressure become dominated by the scalar field potential, assumed to be a constant, and when the kinetic energy of the scalar field becomes negligibly small. Such a dark matter particle production has very important implications on the dynamics and evolution of the Universe. The details of the dark energy-dark matter interaction mechanism depend on the parameters of the particle physics models involved to describe the newly created particles. One of the parameters is the dark matter particle mass, which is a key parameter in the description of the scalar field-dark matter process. Unfortunately presently there is no definite answer giving the value of the mass of the dark matter particle.

Particle creation can be related to what is called the arrow of time: something that provides a direction to time, and distinguishes the future from the past. There are two different arrows of time: the thermodynamical arrow of time, the direction in which entropy increases, and the cosmological arrow of time, the direction in which the Universe is expanding. Particle creation introduces asymmetry in the evolution of the Universe, and enables us to assign a thermodynamical arrow of time, which agrees, in our model, with the cosmological one. This coincidence is determined in a natural way by the decay of the scalar field, due to the presence of a friction force between field and matter.

In our present approach we have neglected the back-reaction of the newly created particles on the dynamics of the Universe, and the effect of the form of the scalar field potential $U(\phi)$ have not been completely envisaged. These aspects of the thermodynamic theory of the dark energy dark matter interaction will be the subject of a future work.

Acknowledgments

FSNL acknowledges financial support of the Fundação para a Ciência e Tecnologia through the grants CERN/FP/123615/2011 and CERN/FP/123618/2011.

[1] A. G. Riess et al., Astron. J. **116**, 1009 (1998); S. Perlmutter et al., Astrophys. J. **517**, 565 (1999); P. de Bernardis et al., Nature **404**, 955 (2000); S. Hanany et al., Astrophys. J. **545**, L5 (2000).

[2] M. Hicken, W. M. Wood-Vasey, S. Blondin, P. Challis, S. Jha, P. L. Kelly, A. Rest, and R. P. Kirshner, Astrophys. J. **700**, 1097 (2009).

[3] E. Komatsu et al. [WMAP Collaboration], Astrophys. J. Suppl. **180**, 330 (2009).

[4] W. J. Percival, S. Cole, D. J. Eisenstein, R. C. Nichol, J. A. Peacock, A. C. Pope and A. S. Szalay, Mon. Not. Roy. Astron. Soc. **381**, 1053 (2007).

[5] P. J. E. Peebles and B. Ratra, Rev. Mod. Phys. **75**, 559 (2003); T. Padmanabhan, Phys. Repts. **380**, 235 (2003).

[6] B. Ratra and P. J. E. Peebles, Phys. Rev **D37**, 3406 (1988); P. J. E. Peebles and B. Ratra, Astrophys. J. Lett. **325**, L17 (1988); R. R. Caldwell, R. Dave, and P. J. Steinhardt, Phys. Rev. Lett. **80**, 1582 (1998).

[7] Y. Fujii and K. Maeda, *The Scalar-Tensor Theory of Gravitation*, Cambridge, Cambridge University Press, (2003); V. Faraoni, *Cosmology in scalar-tensor gravity*, Dordrecht; Boston, Kluwer Academic Publishers, (2004).

[8] P. Salucci, C. Frigerio Martins, and A. Lapi, arXiv:1102.1184 (2011).

[9] J. Binney and S. Tremaine, *Galactic dynamics*, Princeton University Press, Princeton (1987);

[10] M. Persic, P. Salucci, and F. Stel, Month. Not. Roy. Astr. Soc. 281, 27 (1996).

[11] A. Boriello and P. Salucci, Month. Not. Roy. Astr. Soc. **323**, 285 (2001).

[12] J. M. Overduin and P. S. Wesson, Phys. Repts. **402**, 267 (2004).

[13] T. Damour, G. W. Gibbons, and C. C. Gundlach, Phys. Rev. Lett. **64**, 123 (1990); J. A. Casas, J. Garca–Bellido, and M. Quiros, Class. Quant. Grav. **9**, 1371 (1992); C. Wetterich, Astron. Astrophys. **301**, 321 (1995); L. Amendola, Phys. Rev. D **60**, 043501 (1999); L. Amendola, Phys. Rev. D **62**, 043511 (2000).

[14] A. P. Billyard and A. A. Coley, Phys. Rev. D **61**, 083503 (2000); W. Zimdahl and D. Pavon, Phys. Lett. B **521**, 133 (2001); N. Dalal, K. Abazajian, E. E. Jenkins, and A. V. Manohar, Phys. Rev. Lett. **87**, 141302 (2001); L. P. Chimento, A. S. Jakubi, D. Pavon, and W. Zimdahl, Phys. Rev. D **67**, 083513 (2003); G. R. Farrar and P. J. E. Peebles, Astrophys. J. **604**, 1 (2004); R.-G. Cai and A. Wang, JCAP **0503**, 002 (2005); G. Olivares, F. Atrio-Barandela, and D. Pavon, Phys. Rev. D **71**, 063523 (2005); H. M. Sadjadi and M. Alimohammadi, Phys. Rev. D **74**, 103007 (2006); N. J. Nunes and D. F. Mota, Mon. Not. Roy. Astron. Soc. **368**, 751 (2006); Z.-K. Guo, N. Ohta, and S. Tsujikawa, Phys. Rev. D **76**, 023508 (2007) M. Quartin, M. O. Calvao, S. E. Joras, R. R. R. Reis, and I. Waga, JCAP **0805**, 007 (2008); J. Valiviita, E. Majerotto, and R. Maartens, JCAP —bf 0807, 020 (2008); C. G. Boehmer, G. Caldera-Cabral, R. Lazkoz, and R. Maartens, Phys. Rev. D **78**, 023505 (2008); H. Garcia-Compean, G. Garcia-Jimenez, O. Obregon, and C. Ramirez, JCAP **0807**, 016 (2008); T. Barreiro, O. Bertolami, and P. Torres, MNRAS **409** 750 (2010); J.-H. He, B. Wang, and E. Abdalla, Phys. Rev. D **83**, 063515 (2011); L. P. Chimento and M. G. Richarte, Phys. Rev. D **85** 127301 (2012); S. M. R. Micheletti, Phys. Rev. D **85**, 123536 (2012).

[15] F. Cueva Solano and U. Nucamendi, arXiv:1207.0250 [astro-ph.CO] (2012).

[16] C. Kaeonikhom, D. Singleton, S. V. Sushkov, and N. Yongram, arXiv:1209.5219 [gr-qc] (2012).

[17] I. Prigogine, J. Geheniau, E. Gunzig, P. and Nardone, Proc. Natl. Acad. Sci. U.S.A. **85**, 7428.

[18] M. O. Calvao, J. A. S. Lima, and I. Waga, Phys. Lett. A **162**, 223 (1992).

[19] R. Sudharshan and V. B. Johri, 1994, Gen. Rel. Grav. **26**, 41 (1994); J. A. S. Lima, A. S. M. Germano, and L. R. W. Abramo, Phys. Rev. D **53**, 4278 (1996); L. R. W. Abramo, and J. A. S. Lima, J.A.S., Class. Quantum Grav. **13**, 2593 (1996); T. Harko and M. K. Mak, Gen. Rel. Grav. **32**, 865 (2000); G. Steigman, R. C. Santos, and J. A. S. Lima, JCAP **0906**, 033 (2009); A. de Roany and J. A. de Freitas Pacheco, Gen. Rel. Grav. **43**, 61 (2011); J. F. Jesus, F. A. Oliveira, S. Basilakos, and J. A. S. Lima, Phys. Rev. D **84**, 063511 (2011); J. A. S. Lima, S. Basilakos, and F. E. M. Costa, arXiv:1205.0868 (2012).

[20] A. Linde, *Particle Physics and Inflationary Cosmology*, Harwood Academic Publishers (1992); E. W. Kolb and A. Riotto, Phys. Rev. D **55**, 3313 (1997); L. Kofman, A.

Linde, and A. Starobinsky, Phys. Rev. D **56**, 3258 (1997).

[21] J. D. Barrow and P. Saich, Class. Quantum Grav. **10**, 279 (1993).

[22] E. W. Kolb and M. S. Turner, *The Early Universe*, Addison Wesley, Redwood City (1990).